

## CROP : LETTUCE - *BATAVIA*

Trial carried out in France in 2004

### SITE DETAILS

TRIAL N° :	D/F/OIG/03/07	Crop information
10800 VILLY LE MARECHAL North East of France		Variety : Batavia Panthéon Planting date : 25/05/04 Density of planting : 35 cm between plants Cultivation method : Open field Irrigation : over head irrigation

### APPLICATION DETAILS

	Product	Dose rate	Date of spraying	Crop stage at application	Volume of water	Remarks
Application N°1	AMINOFIT.Xtra	5 cc/l	25/05/04	4 to 6 leaves	-	Root soaking till complete moisture of the mini peat clods
Application N°2	AMINOFIT.Xtra	2,5 l/ha	07/06/04	8 to 10 leaves	250 l/ha	9h00, T° = 23°C, H% = 82%, Cloud cover = 0%
Application N°3	AMINOFIT.Xtra	2,5 l/ha	21/06/04	11 days pre harvest	250 l/ha	13h00, T° = 22°C, H% = 86%, Cloud cover = 90%

### MATERIALS AND METHODS

Plot size : 12 plants per plot  
Trial design : 3 repetitions  
Application method : Root soaking and/or foliar spray.  
Equipment used : Hand sprayer equipped with a boom.  
For soaking the roots, the trays full of plants grown in mini peat clods were placed in a vat filled with a solution of water plus AMINOFIT.Xtra till complete moistening of the clods.  
Assessments : Throughout the season : record any visual effect on the plants  
At harvest : weight of 6 lettuces.

### DISCUSSION, CONCLUSION

Climatic conditions were stressful during the progress of the trial with important variations of temperature between day and night.

Once can conclude that :

- One application of AMINOFIT.Xtra to the clods pre transplant followed by 2 foliar applications increase significantly the weight of the lettuces at harvest : + 7.5%,
- Two foliar applications during the crop development increase significantly the weight of the lettuces at harvest : + 6.9 %,
- The pre transplant application by soaking the clods must be completed by foliar applications during the season.

## RESULTS

Date of assessment	02/07/04		
Crop stage	Maturity		
Type of record	Weight of 6 lettuces in g		Average weight per lettuce in g
		% control	
AMINOFIT.Xtra root soaking + 2 foliar sprays	2831 (a)	+ 7.5	472
AMINOFIT.Xtra 2 foliar sprays	2814 (a)	+ 6.9	469
AMINOFIT.Xtra root soaking	2601 (b)	- 1.2	433
UNTREATED CONTROL	2633 (b)	-	439

## GRAPH

**Average weight per lettuce (in g)**

